
Influenza Surveillance Report Week 18: Apr 29th – May 5th, 2018


I. Influenza-like Illness (ILI) Weekly Summary


- Influenza activity is currently low. This risk assessment is based on:
 - The proportion of emergency department visits for ILI was 1.45%, above the regional baseline of 1.06%, and has been decreasing for six weeks.
 - o The proportion of outpatient provider visits for ILI was 0.77%, below the regional baseline of 0.81%, and has been decreasing for one week.
 - The proportion of deaths associated with pneumonia or influenza was 5.73%, below the epidemic threshold of 6.50%.
- One hundred ninety-two (11.54%) laboratory specimens tested positive for influenza: 2 influenza A H3N2, 2 influenza A H1N1, 11 influenza A unknown subtype, and 177 influenza B.
- Two influenza-associated intensive care unit (ICU) hospitalizations were reported during Week 18. Four hundred and seventy-four ICU hospitalizations have been reported since Week 35.
- Since Week 35, 1 influenza-associated pediatric death, 26 clusters of ILI in schools, and 37 influenza outbreaks in long-term care facilities have been reported.
- **Current recommendations**: Encourage respiratory hygiene and <u>prompt treatment</u> <u>with antivirals</u>. Continue to promote vaccination as long as flu viruses are circulating.


Note: Surveillance for ILI in suburban Cook County involves the weekly collection of data from hospitals, physicians' offices, and laboratories. Thank you to all of our surveillance partners for their help in collecting this information.

II. Influenza Activity Level / Disease Burden











III. Circulating Strains³




IV. Seasonal Severity

ICU Hospitalizations⁴


Cumulative Case Count	#	Incidence Rate	Per 100,000
Total Cases	474		20.91
Incidence Rate by Age		Incidence Rate by Region	
0-4	21.64	North	19.51
5-24 25-49	6.61	West	18.85
25-49	4.24	Southwest	23.63
50-64 65+	18.34	South	23.74
65+	92.54		








V. Regional Flu Activity¹



VI. United States Flu Activity⁶



(All data are preliminary and may change as more reports are received.)

For further information about influenza activity see: <u>City of Chicago Surveillance Report</u>, <u>DuPage County Surveillance Report</u>, <u>Illinois Department of Public Health Surveillance Report</u>, <u>CDC Surveillance Report</u>.

Cook County Influenza Surveillance Reports are posted on the <u>CCDPH website</u>. Surveillance data is also viewable in an <u>online, interactive application.</u>

- Data extracted from syndromic surveillance system, ESSENCE; 45 hospital emergency departments (EDs)
 participate in ESSENCE. Influenza-like-illness (ILI) defined as a symptom complex of fever and cough or sore
 throat. ILI = # of ED visits for ILI / total # of ED visits.
- 2. Data extracted from the U.S. Influenza-like Illness Surveillance Network (ILINet); 5 hospitals and 3 physician offices serve as CDC sentinel sites in Cook County. ILI defined as fever ≥ 100°F, cough and/or sore throat in the absence of a known cause other than influenza. ILI = # of visits for ILI / total # of visits.
- 3. Includes viral culture, RT-PCR, and the rapid antigen test. Cases may reside outside suburban Cook County. Participating laboratories: Illinois Department of Public Health Sentinel Laboratories, NorthShore University Health System, Loyola University Medical Center, and ACL Laboratories.
- 4. Cases reported are suburban Cook County residents (excluding Evanston, Skokie, Oak Park, and Stickney) with known age and residence. Cases aggregated by week of admission. Includes all cases reported through the presented week. Rates calculated with 2010 census data.
- 5. Includes all deaths where the immediate cause of death or a contributing factor was pneumonia and/or influenza (aspiration pneumonia excluded). Data includes all of Cook County and has one week lag behind other surveillance indicators. The 3-week running median is displayed. The percentage of deaths due to P&I are compared with a seasonal baseline and epidemic threshold value calculated for each week. Seasonal baseline is calculated using a periodic regression model that incorporates a CDC based robust regression procedure applied to data from the previous four years. An increase of 1.645 standard deviations above the seasonal baseline of P&I deaths is considered the "epidemic threshold," i.e., the point at which the observed proportion of deaths attributed to pneumonia or influenza was significantly higher than would be expected at that time of the year in the absence of substantial influenza-related mortality. http://www.cdc.gov/flu/weekly/overview.htm#Mortality
- 6. Map produced using the proportion of outpatient visits to health care providers for ILI reported through ILINet. Activity levels are compared to the average percent of ILI visits that occur during weeks with little or no influenza virus circulation. http://www.cdc.gov/flu/weekly/index.htm

§Influenza surveillance data are typically aggregated by week. Most years have 52 weeks; however some have 53 weeks. 2014 was a 53-week year. Graphing these years in comparison to 52-week years can be challenging. Because the last week of the calendar year is epidemiologically important for influenza transmission due to holiday family gatherings and school closures, we have graphed these weeks together. In all graphs with 2014-2015 influenza activity represented, Week 53 data appears in the Week 52 column. Consequently, all other data points prior to Week 53 have been moved forward one week, i.e., Week 52 becomes Week 51, Week 51 becomes Week 50, and so on until Week 36 becomes Week 35. This methodology has been adapted from the California Influenza Surveillance Project.